

Multidimensional the Ricci-flat spaces defined by nonlinear equations

Valerii Dryuma

November 14, 2009

- Institute of Mathematics and Computer Science, MD-2028 Chisinau, Republic of Moldova.

1 The Ricci flat 8D metric and KP-equation

In the papers [1]-[2] was showed that the eight-dimensional metric in local coordinates (x, y, z, t, P, Q, U, V)

$$\begin{aligned} {}^8ds^2 = & 2 \left(-PH_{11} - P \frac{\partial F}{\partial t} - H_{12}Q - 2\Gamma_{11}^3 U + H_{22}V \right) dx^2 + \\ & + 4(H_{11}Q - H_{12}V) dx dy + 4 \left(-\frac{\partial F}{\partial z} V + \frac{\partial F}{\partial t} U \right) dx dz + 2(H_{11}U - H_{31}V) dy^2 + \\ & + 2dx dP + 2dy dQ + 2dz dU + 2dt dV, \end{aligned} \quad (1)$$

is the Ricci-flat $R_{ik} = 0$ if the conditions on the functions $H_{ij} = H_{ij}(y, z, t)$, $\Gamma_{11}^3(y, z, t)$ and $F(y, z, t)$

$$\frac{\partial H_{12}}{\partial y} - \frac{\partial H_{22}}{\partial t} = 0, \quad -\frac{\partial H_{11}}{\partial y} + \frac{\partial H_{21}}{\partial t} = 0, \quad -\frac{\partial H_{11}}{\partial z} + \frac{\partial H_{31}}{\partial t} = 0, \quad (2)$$

$$\frac{\partial \Gamma_{11}^3}{\partial z} = 2 \left(\frac{\partial F}{\partial t} \right)^2 + 2H_{11} \frac{\partial F}{\partial t} + 2(H_{11})^2. \quad (3)$$

are hold.

The metric (1) has the form

$$ds^2 = -\Gamma_{jk}^i \xi_i dx^j dx^k + 2d\xi_k dx^k \quad (4)$$

and it is an example of Riemann extension of affinely-connected the four dimensional space in local coordinates x^l with symmetric connection $\Gamma_{jk}^i(x^l) = \Gamma_{kj}^i(x^l)$. From the system (2) after the substitution [3]

$$\begin{aligned} H_{11} = & -\frac{1}{2}u(y, z, t), \quad H_{12} = -\frac{1}{3}v(y, z, t), \quad H_{21} = -\frac{2}{3}v(y, z, t) - \frac{1}{2} \frac{\partial u(y, z, t)}{\partial t} \\ H_{31} = & -\frac{3}{4}w(y, z, t) + \frac{3}{8}u(y, z, t)^2 - \frac{\partial v(y, z, t)}{\partial t} - \frac{1}{2} \frac{\partial^2 u(y, z, t)}{\partial t^2}, \\ H_{22} = & -\frac{1}{2}w(y, z, t) + \frac{1}{2}u(y, z, t)^2 - \frac{\partial v(y, z, t)}{\partial t} - \frac{1}{2} \frac{\partial^2 u(y, z, t)}{\partial t^2} + \frac{\partial u(y, z, t)}{\partial y}, \end{aligned}$$

the famous KP-equation follows

$$\frac{\partial}{\partial t} \left(\frac{\partial u(y, z, t)}{\partial z} - \frac{3}{2} u(y, z, t) \frac{\partial u(y, z, t)}{\partial t} - \frac{1}{4} \frac{\partial^3 u(y, z, t)}{\partial t^3} \right) = \frac{3}{4} \frac{\partial^2 u(y, z, t)}{\partial y^2}. \quad (5)$$

2 4D the Ricci-flat affinely connected subspace and KP-equation

The main result of [2] is the following

Theorem The four-dimensional affinely connected space with non zero coefficients of connection $\Gamma_{jk}^i = \Gamma_{kj}^i$ of the form

$$\Gamma_{11}^1 = H_{11} + \frac{\partial F}{\partial t}, \quad \Gamma_{11}^2 = H_{12}, \quad \Gamma_{11}^3 = \Gamma_{11}^3, \quad \Gamma_{12}^2 = H_{11}, \quad \Gamma_{13}^3 = -\frac{\partial F}{\partial t}$$

$$\Gamma_{22}^3 = -H_{11}, \quad \Gamma_{11}^4 = -H_{22}, \quad \Gamma_{12}^4 = H_{21}, \quad \Gamma_{13}^4 = \frac{\partial F}{\partial z}, \quad \Gamma_{22}^4 = H_{31}$$

is a Ricci flat

$$R_{ij} = \partial_k \Gamma_{ij}^k - \partial_i \Gamma_{kj}^k + \Gamma_{kl}^k \Gamma_{ij}^l - \Gamma_{im}^k \Gamma_{kj}^m = 0,$$

if the conditions (2-3) hold.

The problem of metrizable such type of connection is important in theory of 3-dim manifolds now it is open question.

3 6D the Ricci-flat defined by KP-equation

As particular case we consider the metrics (4) of the form

$$\begin{aligned} {}^6 ds^2 = & \\ = & 4 \left(\frac{\partial}{\partial x} u(x, y, t) \right) P dx dt + 2 dx dP + 4 \left(\frac{\partial}{\partial y} u(x, y, t) \right) P dy dt + 2 dy dQ + \\ + & \left(-2 P u(x, y, t) \frac{\partial}{\partial x} u(x, y, t) - 2 P \frac{\partial^3}{\partial x^3} u(x, y, t) - 2 \mu \left(\frac{\partial}{\partial y} u(x, y, t) \right) Q + 2 \left(\frac{\partial}{\partial x} u(x, y, t) \right) U \right) dt^2 + \\ & + 2 dt dU. \end{aligned} \quad (6)$$

The Ricci tensor such type of the metric

$$\begin{aligned} R_{ij} = & \\ = & \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \left(\frac{\partial}{\partial x} u(x, y, t) \right)^2 + u(x, y, t) \frac{\partial^2}{\partial x^2} u(x, y, t) + \frac{\partial^4}{\partial x^4} u(x, y, t) + \frac{\partial^2}{\partial t \partial x} u(x, y, t) + \mu \frac{\partial^2}{\partial y^2} u(x, y, t) & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \end{aligned}$$

has one component and it is equal to zero on solutions of the KP-equation

$$\frac{\partial(u_t + uu_x + u_{xxx})}{\partial x} + \mu \frac{\partial^2 u}{\partial y^2} = 0.$$

Remark

The metric (3) arises from the Riemann extension of the 3D Einstein-Weyl space

$$^3 ds^2 = dy^2 - 4dxdt - 4u(x, y, t)dt^2, \quad \nu = -4u_x dt,$$

associated with the DKP-equation

$$\frac{\partial(u_t - uu_x)}{\partial x} = \frac{\partial^2 u}{\partial y^2},$$

which is obtained from the Einstein-Weyl conditions on the Ricci tensor

$$R_{ij} + 1/2 \nabla_i \nu_j + 1/4 \nu_i \nu_j - 1/3 (R + 1/2 \nabla^k \nu_k + 1/4 \nu^k \nu_k) h_{ij} = 0$$

4 6D the Ricci-flat metrics associated with KdF-equation

3D-Riemann metric of the form

$$\begin{aligned} ds^2 = & y^2 dx^2 + (l(x, z) y^2 - 1/2) dx dz + 2 dy dz + \\ & + \left((l(x, z))^2 y^2 - 2 \left(\frac{\partial}{\partial x} l(x, z) \right) y + l(x, z) \right) dz^2 \end{aligned} \quad (7)$$

is a flat $R_{ijkl} = 0$ if the function $l(x, z)$ satisfies the KdF-equation (Dryuma, 2006)

$$\frac{\partial^3}{\partial x^3} l(x, z) + \frac{\partial}{\partial z} l(x, z) - 3 l(x, z) \frac{\partial}{\partial x} l(x, z) = 0. \quad (8)$$

It has 13 Christoffel symbols.

Some of them are in form

$$\begin{aligned} \Gamma_{11}^1 = 1/2 \frac{-1 + 2 l(x, z) y^2}{y}, \quad \Gamma_{12}^1 = y^{-1}, \quad \Gamma_{13}^1 = 1/2 \frac{(-1 + 2 l(x, z) y^2) l(x, z)}{y}, \\ \Gamma_{23}^1 = \frac{l(x, z)}{y}, \\ \Gamma_{33}^1 = 1/2 \frac{2 \left(\frac{\partial}{\partial z} l(x, z) \right) y - 4 l(x, z) y \frac{\partial}{\partial x} l(x, z) + 2 \frac{\partial^2}{\partial x^2} l(x, z) - (l(x, z))^2 + 2 (l(x, z))^3 y^2}{y}, \\ \Gamma_{11}^2 = -1/4 \frac{4 y^3 \frac{\partial}{\partial x} l(x, z) - 8 l(x, z) y^2 + 1}{y}, \\ \dots \\ \Gamma_{11}^3 = -y, \quad \Gamma_{13}^3 = -l(x, z) y, \quad \Gamma_{33}^3 = -(l(x, z))^2 y + \frac{\partial}{\partial x} l(x, z). \end{aligned}$$

Six-dimensional Riemann extension of the metrics (4) is the space in local coordinates $(x, y, z, \xi_1, \xi_2, \xi_3)$ and it is defined by the expression

$$ds^2 = -2 \Gamma_{ij}^k \xi_k dx^i dx^j + 2 dx^k \xi_k \quad (9)$$

with a given coefficients Γ_{ij}^k .

Such metric is a flat on solutions of the KdF-equation (8).

To obtain an examples of non a flat $R_{ijkl} \neq 0$ the six-dimensional metrics associated with the KdF-equation it is necessary to introduce an additional terms into the expression (9).

As example the change of the component of metric

$$g_{13} := l(x, z)y^2 - 1/2$$

on the following

$$g_{1,3} := l(x, z)y^2 - 1$$

change radically the Ricci-tensor of the space.

From

$$Matrix(3, 3, (1, 1) = 0, (1, 2) = 0, (1, 3) = 0, (2, 1) = 0, (2, 2) = 0, (2, 3) = 0,$$

$$(3, 1) = 0, (3, 2) = 0, (3, 3) = (-l_{xxx} - l_z + 3l_x)/y,$$

on the

$$Matrix(3, 3, (1, 1) = 0, (1, 2) = 0, (1, 3) = -(1/2)l_x/y, (2, 1) = 0, (2, 2) = 0, (2, 3) = 0,$$

$$(3, 1) = -(1/2)l_x/y, (3, 2) = 0, (3, 3) = (1/2)(-2l_{xxx}y^2 - l_{xx}y - 3l_zy^2 + 7ly^2l_x + l_x)/y^3.$$

In a six-dimensional case it is possible to change components of metric such a way that metric will be Ricci flat $R_{ik} = 0$ on solutions of the KdF-equation but not a flat $R_{ijkl} \neq 0$.

5 6D Heavenly metric and Special Lagrangian equation

We study six-dimensional generalization of the Heavenly metrics

$$\begin{aligned} ds^2 = & dx du + dy dv + dz dw + A(x, y, z, u, v, w) du^2 + \\ & + 2B(x, y, z, u, v, w) du dv + 2E(x, y, z, u, v, w) du dw + C(x, y, z, u, v, w) dv^2 + \\ & + 2H(x, y, z, u, v, w) dv dw + F(x, y, z, u, v, w) dw^2. \end{aligned} \quad (10)$$

The Ricci tensor of the metric (5) has a fifteen components.

Nine of them are equal to zero due the conditions

$$\begin{aligned} \frac{\partial}{\partial u} E(x, y, z, u, v, w) + \frac{\partial}{\partial v} H(x, y, z, u, v, w) + \frac{\partial}{\partial w} F(x, y, z, u, v, w) &= 0, \\ \frac{\partial}{\partial u} B(x, y, z, u, v, w) + \frac{\partial}{\partial v} C(x, y, z, u, v, w) + \frac{\partial}{\partial w} H(x, y, z, u, v, w) &= 0, \\ \frac{\partial}{\partial u} A(x, y, z, u, v, w) + \frac{\partial}{\partial v} B(x, y, z, u, v, w) + \frac{\partial}{\partial w} E(x, y, z, u, v, w) &= 0. \end{aligned} \quad (11)$$

This system of equation has a solutions depending from an arbitrary functions.

In a simplest case we have the solution

$$\begin{aligned}
A(x, y, z, u, v, w) &= \left(\frac{\partial^2}{\partial z^2} f(x, y, z, u, v, w) \right) \frac{\partial^2}{\partial y^2} f(x, y, z, u, v, w) - \left(\frac{\partial^2}{\partial y \partial z} f(x, y, z, u, v, w) \right)^2, \\
C(x, y, z, u, v, w) &= \left(\frac{\partial^2}{\partial x^2} f(x, y, z, u, v, w) \right) \frac{\partial^2}{\partial z^2} f(x, y, z, u, v, w) - \left(\frac{\partial^2}{\partial x \partial z} f(x, y, z, u, v, w) \right)^2, \\
F(x, y, z, u, v, w) &= \left(\frac{\partial^2}{\partial x^2} f(x, y, z, u, v, w) \right) \frac{\partial^2}{\partial y^2} f(x, y, z, u, v, w) - \left(\frac{\partial^2}{\partial x \partial y} f(x, y, z, u, v, w) \right)^2, \\
E(x, y, z, u, v, w) &= \left(\frac{\partial^2}{\partial y \partial z} f(x, y, z, u, v, w) \right) \frac{\partial^2}{\partial x \partial y} f(x, y, z, u, v, w) - \\
&\quad - \left(\frac{\partial^2}{\partial x \partial z} f(x, y, z, u, v, w) \right) \frac{\partial^2}{\partial y^2} f(x, y, z, u, v, w) \\
B(x, y, z, u, v, w) &= \left(\frac{\partial^2}{\partial x \partial z} f(x, y, z, u, v, w) \right) \frac{\partial^2}{\partial y \partial z} f(x, y, z, u, v, w) - \\
&\quad - \left(\frac{\partial^2}{\partial x \partial y} f(x, y, z, u, v, w) \right) \frac{\partial^2}{\partial z^2} f(x, y, z, u, v, w) \\
H(x, y, z, u, v, w) &= \left(\frac{\partial^2}{\partial x \partial z} f(x, y, z, u, v, w) \right) \frac{\partial^2}{\partial x \partial y} f(x, y, z, u, v, w) - \\
&\quad - \left(\frac{\partial^2}{\partial y \partial z} f(x, y, z, u, v, w) \right) \frac{\partial^2}{\partial x^2} f(x, y, z, u, v, w)
\end{aligned}$$

depending from one arbitrary function.

At this conditions the six-dimensional metric looks as

$$\begin{aligned}
{}^6 ds^2 &= \left(\frac{\partial^2}{\partial w^2} K(\vec{x}) \frac{\partial^2}{\partial v^2} K(\vec{x}) - \left(\frac{\partial^2}{\partial v \partial w} K(\vec{x}) \right)^2 \right) dx^2 + \\
&+ 2 \left(\frac{\partial^2}{\partial u \partial w} K(\vec{x}) \frac{\partial^2}{\partial v \partial w} K(\vec{x}) - \frac{\partial^2}{\partial w^2} K(\vec{x}) \frac{\partial^2}{\partial u \partial v} K(\vec{x}) \right) dx dy + \\
&\quad + \left(\frac{\partial^2}{\partial u^2} K(\vec{x}) \frac{\partial^2}{\partial w^2} K(\vec{x}) - \left(\frac{\partial^2}{\partial u \partial w} K(\vec{x}) \right)^2 \right) dy^2 + \\
&+ 2 \left(\frac{\partial^2}{\partial v \partial w} K(\vec{x}) \frac{\partial^2}{\partial u \partial v} K(\vec{x}) - \frac{\partial^2}{\partial u \partial w} K(\vec{x}) \frac{\partial^2}{\partial v^2} K(\vec{x}) \right) dx dz + \\
&\quad + \left(\frac{\partial^2}{\partial v^2} K(\vec{x}) \frac{\partial^2}{\partial u^2} K(\vec{x}) - \left(\frac{\partial^2}{\partial u \partial v} K(\vec{x}) \right)^2 \right) dz^2 + \\
&+ 2 \left(\frac{\partial^2}{\partial u \partial w} K(\vec{x}) \frac{\partial^2}{\partial u \partial v} K(\vec{x}) - \frac{\partial^2}{\partial u^2} K(\vec{x}) \frac{\partial^2}{\partial v \partial w} K(\vec{x}) \right) dz dy + \\
&\quad + dx du + dy dv + dz dw
\end{aligned} \tag{12}$$

where $K(\vec{x}) = K(x, y, z, u, v, w)$ is arbitrary function.

The Ricci tensor R_{ij} of the metric (5) has a six components.

All equations

$$R_{ij} = 0$$

after the substitution

$$K(x, y, z, u, v, w) = \phi(y + v + x, z + w + x)$$

are reduced to the one equation

$$\begin{aligned} & - \left(\frac{\partial^4}{\partial \xi \partial \rho^2 \partial \xi} \phi(\xi, \rho) \right) \frac{\partial^2}{\partial \xi^2} \phi(\xi, \rho) + 2 \left(\frac{\partial^3}{\partial \xi \partial \rho \partial \xi} \phi(\xi, \rho) \right)^2 - 2 \left(\frac{\partial^3}{\partial \rho^2 \partial \xi} \phi(\xi, \rho) \right) \frac{\partial^3}{\partial \xi^3} \phi(\xi, \rho) - \\ & - \left(\frac{\partial^2}{\partial \rho^2} \phi(\xi, \rho) \right) \frac{\partial^4}{\partial \xi^4} \phi(\xi, \rho) + 2 \left(\frac{\partial^2}{\partial \rho \partial \xi} \phi(\xi, \rho) \right) \frac{\partial^4}{\partial \xi^2 \partial \rho \partial \xi} \phi(\xi, \rho) + \\ & + 2 \left(\frac{\partial^3}{\partial \rho^2 \partial \xi} \phi(\xi, \rho) \right)^2 - \left(\frac{\partial^4}{\partial \rho^4} \phi(\xi, \rho) \right) \frac{\partial^2}{\partial \xi^2} \phi(\xi, \rho) - 2 \left(\frac{\partial^3}{\partial \rho^3} \phi(\xi, \rho) \right) \frac{\partial^3}{\partial \xi \partial \rho \partial \xi} \phi(\xi, \rho) - \\ & - \left(\frac{\partial^2}{\partial \rho^2} \phi(\xi, \rho) \right) \frac{\partial^4}{\partial \xi \partial \rho^2 \partial \xi} \phi(\xi, \rho) + 2 \left(\frac{\partial^2}{\partial \rho \partial \xi} \phi(\xi, \rho) \right) \frac{\partial^4}{\partial \rho^3 \partial \xi} \phi(\xi, \rho) = 0, \quad (13) \end{aligned}$$

where

$$\xi = x + y + v, \quad \rho = z + x + w.$$

In compact form this equation can be rewritten as

$$\Delta \psi(\xi, \rho) = 0$$

where

$$\psi(\xi, \rho) = \left(\frac{\partial^2}{\partial \xi^2} \phi(\xi, \rho) \right) \frac{\partial^2}{\partial \rho^2} \phi(\xi, \rho) - \left(\frac{\partial^2}{\partial \xi \partial \rho} \phi(\xi, \rho) \right)^2$$

and

$$\Delta = \frac{\partial^2}{\partial \xi^2} + \frac{\partial^2}{\partial \rho^2}$$

is the Laplace operator.

Its solutions give the Ricci-flat examples of the metric (5).

5.1 The Beltrami parameters

To the investigation of the properties of the metrics (5) can be considered two invariant equations defined by the first

$$\Delta \psi = g^{ij} \frac{\partial \psi}{\partial x^i} \frac{\partial \psi}{\partial x^j}$$

and the second

$$\square \psi = g^{ij} \left(\frac{\partial^2}{\partial x^i \partial x^j} - \Gamma_{ij}^k \frac{\partial}{\partial x^k} \right) \psi$$

Beltrami parameters.

To the metric (5) the equation $\square \phi = 0$ looks as ($K = \phi(\vec{x})$)

$$\frac{\partial^2}{\partial u \partial x} \phi(\vec{x}) + \frac{\partial^2}{\partial w \partial z} \phi(\vec{x}) + \frac{\partial^2}{\partial v \partial y} \phi(\vec{x}) - \left(\frac{\partial^2}{\partial y^2} \phi(\vec{x}) \right) \left(\frac{\partial^2}{\partial x^2} \phi(\vec{x}) \right) \frac{\partial^2}{\partial z^2} \phi(\vec{x}) +$$

$$\begin{aligned}
& +2 \left(\frac{\partial^2}{\partial x \partial z} \phi(\vec{x}) \right) \left(\frac{\partial^2}{\partial x \partial z} f(\vec{x}) \right) \frac{\partial^2}{\partial y^2} f(\vec{x}) - 2 \left(\frac{\partial^2}{\partial x \partial z} \phi(\vec{x}) \right) \left(\frac{\partial^2}{\partial y \partial z} f(\vec{x}) \right) \frac{\partial^2}{\partial x \partial y} f(\vec{x}) + \\
& +2 \left(\frac{\partial^2}{\partial x \partial y} \phi(\vec{x}) \right) \left(\frac{\partial^2}{\partial x \partial y} f(\vec{x}) \right) \frac{\partial^2}{\partial z^2} f(\vec{x}) - \left(\frac{\partial^2}{\partial x^2} \phi(\vec{x}) \right) \left(\frac{\partial^2}{\partial z^2} f(\vec{x}) \right) \frac{\partial^2}{\partial y^2} f(\vec{x}) + \\
& +2 \left(\frac{\partial^2}{\partial y \partial z} \phi(\vec{x}) \right) \left(\frac{\partial^2}{\partial y \partial z} f(\vec{x}) \right) \frac{\partial^2}{\partial x^2} f(\vec{x}) - 2 \left(\frac{\partial^2}{\partial y \partial z} \phi(\vec{x}) \right) \left(\frac{\partial^2}{\partial x \partial z} f(\vec{x}) \right) \frac{\partial^2}{\partial x \partial y} f(\vec{x}) - \\
& - \left(\frac{\partial^2}{\partial z^2} \phi(\vec{x}) \right) \left(\frac{\partial^2}{\partial x^2} f(\vec{x}) \right) \frac{\partial^2}{\partial y^2} f(\vec{x}) - 2 \left(\frac{\partial^2}{\partial x \partial y} \phi(\vec{x}) \right) \left(\frac{\partial^2}{\partial x \partial z} f(\vec{x}) \right) \frac{\partial^2}{\partial y \partial z} f(\vec{x}) + \\
& + \left(\frac{\partial^2}{\partial z^2} \phi(\vec{x}) \right) \left(\frac{\partial^2}{\partial x \partial y} f(\vec{x}) \right)^2 + \left(\frac{\partial^2}{\partial y^2} \phi(\vec{x}) \right) \left(\frac{\partial^2}{\partial x \partial z} f(\vec{x}) \right)^2 + \\
& + \left(\frac{\partial^2}{\partial x^2} \phi(\vec{x}) \right) \left(\frac{\partial^2}{\partial y \partial z} f(\vec{x}) \right)^2 = 0. \tag{14}
\end{aligned}$$

In particular case the equation (5.1) after the substitution

$$\phi(\vec{x}) = f(\vec{x})$$

takes the form

$$\begin{aligned}
& \frac{\partial^2}{\partial u \partial x} f(\vec{x}) + \frac{\partial^2}{\partial w \partial z} f(\vec{x}) + \frac{\partial^2}{\partial v \partial y} f(\vec{x}) - 3 \left(\frac{\partial^2}{\partial y^2} f(\vec{x}) \right) \left(\frac{\partial^2}{\partial x^2} f(\vec{x}) \right) \frac{\partial^2}{\partial z^2} f(\vec{x}) + \\
& + 3 \left(\frac{\partial^2}{\partial x \partial z} f(\vec{x}) \right)^2 \frac{\partial^2}{\partial y^2} f(\vec{x}) - 6 \left(\frac{\partial^2}{\partial x \partial z} f(\vec{x}) \right) \left(\frac{\partial^2}{\partial y \partial z} f(\vec{x}) \right) \frac{\partial^2}{\partial x \partial y} f(\vec{x}) + \\
& + 3 \left(\frac{\partial^2}{\partial x \partial y} f(\vec{x}) \right)^2 \frac{\partial^2}{\partial z^2} f(\vec{x}) + 3 \left(\frac{\partial^2}{\partial y \partial z} f(\vec{x}) \right)^2 \frac{\partial^2}{\partial x^2} f(\vec{x}) = 0. \tag{15}
\end{aligned}$$

After the change of variables

$$f(\vec{x}) = f(x, y, z, u, v, w) = h(x + u, v + y, w + z) = h(\eta, \xi, \rho)$$

the equation (5.1) is reduced to the form

$$\begin{aligned}
& \frac{\partial^2}{\partial \eta^2} h(\eta, \xi, \rho) + \frac{\partial^2}{\partial \rho^2} h(\eta, \xi, \rho) + \frac{\partial^2}{\partial \xi^2} h(\eta, \xi, \rho) + 3 \left(\frac{\partial^2}{\partial \eta \partial \rho} h(\eta, \xi, \rho) \right)^2 \frac{\partial^2}{\partial \xi^2} h(\eta, \xi, \rho) - \\
& - 6 \left(\frac{\partial^2}{\partial \eta \partial \rho} h(\eta, \xi, \rho) \right) \left(\frac{\partial^2}{\partial \rho \partial \xi} h(\eta, \xi, \rho) \right) \frac{\partial^2}{\partial \eta \partial \xi} h(\eta, \xi, \rho) + 3 \left(\frac{\partial^2}{\partial \eta \partial \xi} h(\eta, \xi, \rho) \right)^2 \frac{\partial^2}{\partial \rho^2} h(\eta, \xi, \rho) - \\
& - 3 \left(\frac{\partial^2}{\partial \xi^2} h(\eta, \xi, \rho) \right) \left(\frac{\partial^2}{\partial \eta^2} h(\eta, \xi, \rho) \right) \frac{\partial^2}{\partial \rho^2} h(\eta, \xi, \rho) + 3 \left(\frac{\partial^2}{\partial \rho \partial \xi} h(\eta, \xi, \rho) \right)^2 \frac{\partial^2}{\partial \eta^2} h(\eta, \xi, \rho) = 0 \tag{16}
\end{aligned}$$

or

$$\Delta h(\eta, \xi, \rho) - 3 \det \begin{bmatrix} \frac{\partial^2}{\partial \eta^2} h(\eta, \xi, \rho) & \frac{\partial^2}{\partial \eta \partial \xi} h(\eta, \xi, \rho) & \frac{\partial^2}{\partial \eta \partial \rho} h(\eta, \xi, \rho) \\ \frac{\partial^2}{\partial \eta \partial \xi} h(\eta, \xi, \rho) & \frac{\partial^2}{\partial \xi^2} h(\eta, \xi, \rho) & \frac{\partial^2}{\partial \rho \partial \xi} h(\eta, \xi, \rho) \\ \frac{\partial^2}{\partial \eta \partial \rho} h(\eta, \xi, \rho) & \frac{\partial^2}{\partial \rho \partial \xi} h(\eta, \xi, \rho) & \frac{\partial^2}{\partial \rho^2} h(\eta, \xi, \rho) \end{bmatrix} = 0,$$

where

$$\Delta = \frac{\partial^2}{\partial \eta^2} + \frac{\partial^2}{\partial \xi^2} + \frac{\partial^2}{\partial \rho^2}$$

is a three-dimensional Laplace operator.

The equation (5.1) is a famous Harvey-Lawson "Special Lagrangian" equation having an important applications in theory of Calabi-Yau manifolds and mirror symmetry.

5.2 The simplest solutions

To obtain particular solutions of the partial nonlinear differential equation

$$F(x, y, z, f_x, f_y, f_z, f_{xx}, f_{xy}, f_{xz}, f_{yy}, f_{yz}, f_{xxx}, f_{xyy}, f_{xxy}, \dots) = 0 \quad (17)$$

can be applied a following approach.

We use the following parametric presentation of the functions and variables

$$\begin{aligned} f(x, y, z) &\rightarrow u(x, t, z), \quad y \rightarrow v(x, t, z), \quad f_x \rightarrow u_x - \frac{u_t}{v_t} v_x, \\ f_z &\rightarrow u_z - \frac{u_t}{v_t} v_z, \quad f_y \rightarrow \frac{u_t}{v_t}, \quad f_{yy} \rightarrow \frac{(\frac{u_t}{v_t})_t}{v_t}, \quad f_{xy} \rightarrow \frac{(u_x - \frac{u_t}{v_t} v_x)_t}{v_t}, \dots \end{aligned} \quad (18)$$

where variable t is considered as parameter.

Remark that conditions of the type

$$f_{xy} = f_{yx}, \quad f_{xz} = f_{zx} \dots$$

hold at the such type of presentation.

In result instead of equation (17) one get the relation between the new variables $u(x, t, z)$ and $v(x, t, z)$ and their partial derivatives

$$\Psi(u, v, u_x, u_z, u_t, v_x, v_z, v_t, \dots) = 0. \quad (19)$$

This relation coincides with initial p.d.e at the condition $v(x, t, z) = t$ and takes more general form after presentation of the functions u, v in form $u(x, t, z) = F(\omega, \omega_t \dots)$ and $v(x, t, z, s) = \Phi(\omega, \omega_t \dots)$ with some function $\omega(x, t, z)$.

In result of change variables and function with accordance of (5.2) the equation (5.1)

$$\begin{aligned} &\frac{\partial^2}{\partial x^2} h(x, y, z) + \frac{\partial^2}{\partial z^2} h(x, y, z) + \frac{\partial^2}{\partial y^2} h(x, y, z) + 3 \left(\frac{\partial^2}{\partial x \partial z} h(x, y, z) \right)^2 \frac{\partial^2}{\partial y^2} h(x, y, z) - \\ &- 6 \left(\frac{\partial^2}{\partial x \partial z} h(x, y, z) \right) \left(\frac{\partial^2}{\partial y \partial z} h(x, y, z) \right) \frac{\partial^2}{\partial x \partial y} h(x, y, z) + 3 \left(\frac{\partial^2}{\partial x \partial y} h(x, y, z) \right)^2 \frac{\partial^2}{\partial z^2} h(x, y, z) - \\ &- 3 \left(\frac{\partial^2}{\partial y^2} h(x, y, z) \right) \left(\frac{\partial^2}{\partial x^2} h(x, y, z) \right) \frac{\partial^2}{\partial z^2} h(x, y, z) + 3 \left(\frac{\partial^2}{\partial y \partial z} h(x, y, z) \right)^2 \frac{\partial^2}{\partial x^2} h(x, y, z) = 0 \end{aligned}$$

is transformed into the relation (19).

This relation after the substitution

$$u(x, t, z) = t \frac{\partial}{\partial t} \omega(x, t, z) - \omega(x, t, z), \quad v(x, t, z) = \frac{\partial}{\partial t} \omega(x, t, z). \quad (20)$$

takes the form of p.d.e.

$$\begin{aligned}
& -3 \left(\frac{\partial^2}{\partial x^2} \omega(x, t, z) \right) \frac{\partial^2}{\partial z^2} \omega(x, t, z) + 3 \left(\frac{\partial^2}{\partial x \partial z} \omega(x, t, z) \right)^2 - \left(\frac{\partial^2}{\partial t^2} \omega(x, t, z) \right) \frac{\partial^2}{\partial x^2} \omega(x, t, z) + \\
& + 1 + \left(\frac{\partial^2}{\partial t \partial z} \omega(x, t, z) \right)^2 + \left(\frac{\partial^2}{\partial t \partial x} \omega(x, t, z) \right)^2 - \left(\frac{\partial^2}{\partial t^2} \omega(x, t, z) \right) \frac{\partial^2}{\partial z^2} \omega(x, t, z) = 0.
\end{aligned} \tag{21}$$

The equation (5.2) consists from the 2D M-A equations with respect the variables (x, t) , (x, z) and (t, z) .

Let us consider some examples of its solutions.

1. To the equation

$$\mu \Delta(f) + Hess(f) = 0 \tag{22}$$

The substitution into (22)

$$\omega(x, t, z) = A(x^2 + z^2, t)$$

lead to the equation on the function $A(\eta, t)$, ($\eta = x^2 + z^2$)

$$\begin{aligned}
& -4\mu \left(\frac{\partial^2}{\partial \eta \partial t} A(\eta, t) \right)^2 \eta - 8 \left(\frac{\partial^2}{\partial \eta^2} A(\eta, t) \right) \eta \frac{\partial}{\partial \eta} A(\eta, t) - 4 \left(\frac{\partial}{\partial \eta} A(\eta, t) \right)^2 + \\
& + 4\mu \left(\frac{\partial^2}{\partial t^2} A(\eta, t) \right) \frac{\partial}{\partial \eta} A(\eta, t) - \mu + 4\mu \left(\frac{\partial^2}{\partial t^2} A(\eta, t) \right) \frac{\partial^2}{\partial \eta^2} A(\eta, t) = 0.
\end{aligned}$$

Its particular solution is

$$A(\eta, t) = B(t) + \eta e^{kt},$$

$$B(t) = \frac{e^{kt}}{\mu k^2} + 1/4 \frac{e^{-kt}}{k^2} + C_1 t + C_2.$$

Now elimination of the parameter t from the system

$$y - \frac{\partial}{\partial t} \omega(x, t, z) = 0, \quad f(x, y, z) - t \frac{\partial}{\partial t} \omega(x, t, z) + \omega(x, t, z) = 0$$

lead to the solution of the equation (22)

$$\begin{aligned}
f(x, y, z) = & - \left(1 - \ln \left(\frac{y\mu + \sqrt{T}}{1 + \mu x^2 + \mu z^2} \right) y\sqrt{T} + \ln(2) y^2 \mu - \ln \left(\frac{y\mu + \sqrt{T}}{1 + \mu x^2 + \mu z^2} \right) y^2 \mu \right) (y\mu + \sqrt{T})^{-1} + \\
& + (y^2 \mu + \mu x^2) + \left(\ln(2) y\sqrt{T} + \mu z^2 + y\sqrt{T} \right) (y\mu + \sqrt{T})^{-1},
\end{aligned}$$

where

$$\mu (y^2 \mu + 1 + \mu x^2 + \mu z^2) = T$$

2. The substitution

$$\omega(x, t, z) = A(x + t, z)$$

into (22) lead to the equation on the function $A(\eta, z)$ ($\eta = x + z$)

$$- \left(\frac{\partial^2}{\partial \eta^2} A(\eta, z) \right) \frac{\partial^2}{\partial z^2} A(\eta, z) - \mu \left(\frac{\partial^2}{\partial \eta \partial z} A(\eta, z) \right)^2 + \mu \left(\frac{\partial^2}{\partial \eta^2} A(\eta, z) \right) \frac{\partial^2}{\partial z^2} A(\eta, z) -$$

$$-\mu + \left(\frac{\partial^2}{\partial \eta \partial z} A(\eta, z) \right)^2$$

At the $\mu = -1/3$ we get the M-A equation

$$-\left(\frac{\partial^2}{\partial \eta^2} A(\eta, z) \right) \frac{\partial^2}{\partial z^2} A(\eta, z) + \left(\frac{\partial^2}{\partial \eta \partial z} A(\eta, z) \right)^2 + 1/4 = 0 \quad (23)$$

which after the (u, v) -transformation is reduced to the Laplace equation

$$4 \frac{\partial^2}{\partial \eta^2} \theta(\eta, \xi) + \frac{\partial^2}{\partial \xi^2} \theta(\eta, \xi) = 0.$$

Its general solution

$$\theta(\eta, \xi) = M(\eta + 2I\xi) + N(\eta - 2I\xi)$$

allow us to construct particular solutions of the equation (23) and corresponding “Special lagrangian equaion” .

6 Heavenly metrics and Yang-Mills equation

We consider following generalization of Heavenly metrics

$$\begin{aligned} ds^2 = & dx du + dy dv + dz dw + A(x, y, z, p, q) du^2 + 2 B(x, y, z, p, q) du dv + \\ & + 2 E(x, y, z, p, q) du dw + C(x, y, z, p, q) dv^2 + 2 H(x, y, z, p, q) dv dw + \\ & + F(x, y, z, p, q) dw^2 + dp dq, \end{aligned} \quad (24)$$

having a following components

$$\begin{aligned} A(x, y, z, p, q) &= \left(\frac{\partial^2}{\partial z^2} f(x, y, z, p, q) \right) \frac{\partial^2}{\partial y^2} f(x, y, z, p, q) - \left(\frac{\partial^2}{\partial y \partial z} f(x, y, z, p, q) \right)^2, \\ C(x, y, z, p, q) &= \left(\frac{\partial^2}{\partial x^2} f(x, y, z, p, q) \right) \frac{\partial^2}{\partial z^2} f(x, y, z, p, q) - \left(\frac{\partial^2}{\partial x \partial z} f(x, y, z, p, q) \right)^2, \\ F(x, y, z, p, q) &= \left(\frac{\partial^2}{\partial x^2} f(x, y, z, p, q) \right) \frac{\partial^2}{\partial y^2} f(x, y, z, p, q) - \left(\frac{\partial^2}{\partial x \partial y} f(x, y, z, p, q) \right)^2, \\ E(x, y, z, p, q) &= \left(\frac{\partial^2}{\partial y \partial z} f(x, y, z, p, q) \right) \frac{\partial^2}{\partial x \partial y} f(x, y, z, p, q) - \left(\frac{\partial^2}{\partial x \partial z} f(x, y, z, p, q) \right) \frac{\partial^2}{\partial y^2} f(x, y, z, p, q), \\ B(x, y, z, p, q) &= \left(\frac{\partial^2}{\partial x \partial z} f(x, y, z, p, q) \right) \frac{\partial^2}{\partial y \partial z} f(x, y, z, p, q) - \left(\frac{\partial^2}{\partial x \partial y} f(x, y, z, p, q) \right) \frac{\partial^2}{\partial z^2} f(x, y, z, p, q), \\ H(x, y, z, p, q) &= \left(\frac{\partial^2}{\partial x \partial z} f(x, y, z, p, q) \right) \frac{\partial^2}{\partial x \partial y} f(x, y, z, p, q) - \left(\frac{\partial^2}{\partial y \partial z} f(x, y, z, p, q) \right) \frac{\partial^2}{\partial x^2} f(x, y, z, p, q). \end{aligned}$$

At these conditions from 21 components of the Ricci-tensor of the metric (6) only six components

$$R_{uu} \neq 0, \quad R_{uv} \neq 0, \quad R_{uw} \neq 0, \quad R_{vv} \neq 0, \quad R_{vw} \neq 0, \quad R_{ww} \neq 0.$$

are different from zero.

The equation

$$g^{ij} \left(\frac{\partial^2 \Psi}{\partial x^i \partial x^j} - \Gamma_{ij}^k \frac{\partial \Psi}{\partial x^k} \right) = 0,$$

defined by the Laplace-Beltrami operator of the metric (6) depends on the function $f(x, y, z, p, q)$ and after the substitution $f = \Psi$ takes the form

$$\begin{aligned} & \frac{\partial^2}{\partial p \partial q} f(x, y, z, p, q) + \\ & + 3 \left(\frac{\partial^2}{\partial x \partial y} f(x, y, z, p, q) \right)^2 \frac{\partial^2}{\partial z^2} f(x, y, z, p, q) - \\ & - 6 \left(\frac{\partial^2}{\partial x \partial y} f(x, y, z, p, q) \right) \left(\frac{\partial^2}{\partial x \partial z} f(x, y, z, p, q) \right) \frac{\partial^2}{\partial y \partial z} f(x, y, z, p, q) + \\ & + 3 \left(\frac{\partial^2}{\partial y \partial z} f(x, y, z, p, q) \right)^2 \frac{\partial^2}{\partial x^2} f(x, y, z, p, q) - \\ & - 3 \left(\frac{\partial^2}{\partial y^2} f(x, y, z, p, q) \right) \left(\frac{\partial^2}{\partial x^2} f(x, y, z, p, q) \right) \frac{\partial^2}{\partial z^2} f(x, y, z, p, q) + \\ & + 3 \left(\frac{\partial^2}{\partial x \partial z} f(x, y, z, p, q) \right)^2 \frac{\partial^2}{\partial y^2} f(x, y, z, p, q) = 0 \end{aligned}$$

or

$$\begin{aligned} & \frac{\partial^2}{\partial p \partial q} f(x, y, z, p, q) - \\ & - 3 \left[\begin{array}{ccc} \frac{\partial^2}{\partial x^2} f(x, y, z, p, q) & \frac{\partial^2}{\partial x \partial y} f(x, y, z, p, q) & \frac{\partial^2}{\partial x \partial z} f(x, y, z, p, q) \\ \frac{\partial^2}{\partial x \partial y} f(x, y, z, p, q) & \frac{\partial^2}{\partial y^2} f(x, y, z, p, q) & \frac{\partial^2}{\partial y \partial z} f(x, y, z, p, q) \\ \frac{\partial^2}{\partial x \partial z} f(x, y, z, p, q) & \frac{\partial^2}{\partial y \partial z} f(x, y, z, p, q) & \frac{\partial^2}{\partial z^2} f(x, y, z, p, q) \end{array} \right] = 0. \quad (25) \end{aligned}$$

The equation (6) is of Monge-Ampere type and after the substitution

$$f(x, y, z, p, q) = U(p, q)E(x, y, z)$$

takes the form

$$\frac{\frac{\partial^2}{\partial p \partial q} U(p, q)}{(U(p, q))^3} - 3 \frac{Hess(E(x, y, z))}{E(x, y, z)} = 0.$$

So it has particular solutions defined by the equations

$$3Hess(E) - \mu E = 0$$

and

$$\frac{\partial^2 U}{\partial p \partial q} - \mu U^3 = 0.$$

Both of equations have an important applications.

The second equation is reduction of the $SU(2)$ Yang-Mills equation and the first is a Monge-Ampere type of equation having applications in theory of Calabi-Yau manifolds.

References

- [1] V.Dryuma: Eight-dimensional the Ricci-flat Riemann space related with the KP-equation. *arXiv:0810.0346 v1 2 Oct 2008*, 1–5, *v2 15 Oct 2008* 1–5.
- [2] V.Dryuma: Four-dimensional the Ricci-flat space defined by the KP-equation. *Buletinul Academiei de Stiinte a Republicii Moldova*, **3** (2008) 108–111.
- [3] B. Konopelchenko: Quantum deformations of associative algebras and integrable systems. *arXiv:0802.3022 v2 [nlin.SI] 28 Apr 2008* 1–24.
- [4] V.Dryuma: The Riemann and Einstein-Weyl geometries in the theory of ODE's, their applicatons and all that. *In: New Trends in Integrability and Partial Solvability (NATO Sci. ser.II: Math., Phys., Chem., vol.132, A.B.Shabat, A. Conzales-Lopez, M.Manas, L.Martinez Alonso and M.A.Rodrigues)*, Kluwe, Dordrecht (2004), pp.115-156.
- [5] V.Dryuma: Toward a theory of spaces of constant curvature. *Theoretical an Mathematical Physics*, 146(1): 35-45 (2006)
- [6] V.Dryuma, L. Bogdanov : On nonlinear equations connected with six-dimensional Plebanski space. *arXiv:0812.1637 v1 [physics.gen-ph] 9 Dec 2008* 1–11.